

Generic Name: fidaxomicin

Applicable Drugs: Dificid®

Preferred: N/A

Non-preferred: N/A

Date of Origin: 12/13/2023

Date Last Reviewed / Revised: N/A

PRIOR AUTHORIZATION CRITERIA

(May be considered medically necessary when criteria I through V are met)

- I. Documented presence of clinically significant diarrhea caused by *Clostridioides difficile* infection that is confirmed by positive stool assay.
- II. Age \geq 6 months.
- III. Documentation of treatment failure or contraindication to a course of oral vancomycin.
- IV. Medication is prescribed in accordance with FDA labeling and is supported by current clinical practice guidelines.
- V. Treatment must be prescribed by or in consultation with a gastroenterologist or infectious disease specialist.

EXCLUSION CRITERIA

- N/A

OTHER CRITERIA

- N/A

QUANTITY / DAYS SUPPLY RESTRICTIONS

- Adults and pediatric patients weighing at least 12.5 kg: twenty 200 mg tablets per 10 days.
 - Patients weighing more than 12.5 kg who cannot swallow pills may be approved for one bottle of 40 mg/mL oral suspension per 10 days instead.
- Pediatric patients weighing at least 4 kg: one bottle of 40 mg/mL oral suspension per 10 days.

APPROVAL LENGTH

- **Authorization:** One course of treatment (10 days).
- **Re-Authorization:** N/A

APPENDIX

REFERENCES

1. Dificid. Prescribing Information. Merck & Co., Inc.; 2022. Accessed December 13, 2023. https://www.merck.com/product/usa/pi_circulars/d/dificid/dificid_pi.pdf
2. Infectious Diseases Society of America and Society for Healthcare Epidemiology of America. 2021 Focused Update Guidelines on Management of Clostridioides difficile Infection in Adults. Accessed December 13, 2023. <https://www.idsociety.org/practice-guideline/clostridioides-difficile-2021-focused-update/>

DISCLAIMER: Medication Policies are developed to help ensure safe, effective and appropriate use of selected medications. They offer a guide to coverage and are not intended to dictate to providers how to practice medicine. Refer to Plan for individual adoption of specific Medication Policies. Providers are expected to exercise their medical judgement in providing the most appropriate care for their patients.